

**Intensive critical care and management of asthmatic and smoker patients
in COVID-19 infection**

DONGMING LU¹

OBAID YAQOUB²

MANISH KUMAR^{3,4}

AJAY SINGH KUSHWAH^{2,*}

RAHUL KUMAR SHARMA²

DEVINDER KUMAR²

YOGENDRA MAVAI⁵

RUKAIYA KHAN⁶

¹ *Department of Pulmonary and Critical Care Medicine, Jinling Hospital
Nanjing University School of Medicine, Nanjing, Jiangsu, 210000, China*

² *Department of Pharmacology, Amar Shaheed Baba Ajit Singh Jujhar Singh Memorial College of
Pharmacy, Bela (An Autonomous College), Ropar, Punjab, 140111, India*

³ *Chitkara College of Pharmacy, Chitkara University, Punjab, 140401, India*

⁴ *Department of Neurosurgery, College of Medicine, Penn State Health Milton S. Hershey Medical
Center, The Pennsylvania State University, Pennsylvania 17033-0850, USA*

⁵ *IPS College of Pharmacy, Shiypuri Link Road Gwalior, M.P., 474001, India*

⁶ *Abhilashi College of Pharmacy, Nerchowk Mandi, H.P., 175008, India*

*Correspondence; e-mail: kushwah_ph05@yahoo.co.in; ajay@copbela.org

ABSTRACT

This century's most serious catastrophe, COVID-19, has been dubbed "the most life-threatening disaster ever". Asthmatic persons are even more prone to COVID-19's complex interplay with the underlying inflammatory condition. In order to protect themselves against COVID-19, asthmatic patients must be

31 very vigilant in their usage of therapeutic techniques and drugs (*e.g.*, bronchodilators, 5-lipoxygenase
32 inhibitors), which may be accessed to deal with mild, moderate, and severe COVID-19 indications.
33 People with asthma may have more severe COVID-19 symptoms, which may lead to a worsening of
34 their condition. Several cytokines were found to be elevated in the bronchial tracts of patients with acute
35 instances of COVID-19, suggesting that this ailment may aggravate asthma episodes by increasing
36 inflammation. The intensity of COVID-19 symptoms is lessened in patients with asthma who have
37 superior levels of T-cells. Several antibiotics, antivirals, antipyretics, and anti-inflammatory drugs have
38 been suggested to suppress COVID-19 symptoms in asthmatic persons. Furthermore, smokers are more
39 likely to have aggravated repercussions in COVID-19 infection. Being hospitalized to critical care due
40 to COVID-19, needing mechanical breathing, and suffering from serious health repercussions, are all
41 possible outcomes for someone who has previously smoked. Smoking damages airways and alveoli,
42 which significantly raises the risk of COVID-19-related health complications. Patients with a previous
43 record of smoking are predisposed to severe COVID-19 disease symptoms that essentially require a
44 combination of bronchodilators, mucolytics, antivirals, and antimuscarinic drugs, to cope with the
45 situation. The present review discusses the care and management of asthmatic and smoker patients in
46 COVID-19 infection.

47 *Keywords:* COVID-19, asthma, smoking, critical care, SARS-CoV-2

48

49

AN OVERVIEW OF COVID-19

50 Coronavirus is a genus of viruses that may infect people, birds, and animals, producing colds,
51 coughs, fevers, and alveolar pneumonia-like symptoms, such as shortness of breath. COVID-
52 19 is hailed as the utmost severe epidemic of the 21st century. It was widely assumed at the onset
53 that it began at a seafood market in Wuhan, China (1). The World Health Organization (WHO)
54 designated this virus the 2019-nCov, which is a scientific word meaning "new" (n for a novel).
55 Bats are thought to be the virus's natural reservoir since traces of the virus were detected in bats
56 for the first time. SARS-CoV-2 (Severe Acute Respiratory Syndrome Coronavirus 2), COVID-
57 19, the 2019 Novel Coronavirus (2019-nCov), the 2019 Coronavirus, and the Wuhan
58 Coronavirus are all names for this neoteric virus. Because of the high rate of transmission,
59 people are more susceptible to this viral illness. Fever, tiredness, dry cough, sore throat, and
60 breathing problems are common symptoms in COVID-19 patients (2). On January 30, 2020,
61 COVID-19 has asserted a public health crisis, and on March 11, 2020, it was announced a
62 worldwide pandemic (3). The most impacted countries were the United States, India, France,
63 Germany, United Kingdom, Italy, China and Brazil. In July 2022, approx. 580 million total
64 cases of Covid-19 worldwide were recorded, and still, there are about 23.5 million total active

65 cases as of July 31, 2022. Approximately 550 million people have recovered, yet about 6.4
66 million people have died. COVID-19 positive cases were confirmed in about 44 million persons
67 in India, with a recovery rate of around 98 %. This virus has a 5.1-day incubation period on
68 average (2-14 days). COVID-19 symptoms include fever, lethargy, dry cough, muscular
69 discomfort, dyspnea, pneumonia, and, in many instances, a reduction in leukocyte counts. In
70 addition to the aching (tender) pharynx, diarrhea, pink eye (inflammation of the conjunctiva),
71 headache, skin rashes, and discoloration of the fingers, digits, or toes, COVID-19 symptoms
72 include high temperature, chills and tiredness (4). COVID-19 was first thought to be a
73 widespread respiratory illness in 2020, however, it was subsequently discovered that persons
74 with asthma and smokers would be at an elevated risk of severe disease and death (5). Smoking
75 and using E-cigarettes both increase the likelihood and severity of lung infections, owing to the
76 damage done to the upper airways and the resulting loss in pulmonary immune function.
77 Smokers, in particular, are at an elevated hazard of illness and death owing to coronavirus
78 infection (6). We focus on the link between COVID-19 and asthma/smoking, as well as their
79 therapeutic treatment, in this review.

80

81

ASTHMA AND COVID-19

82 *Hazards of COVID-19 to asthmatic patients*

83 People of all ages may suffer from asthma, including children, adults and the elderly.
84 Asthma is defined by a rise in hyper-responsiveness to the tracheobronchial structures caused
85 by a variety of peripheral provocations, which finally narrow the airway tract. Asthma is
86 described as "a long-term inflammatory condition that aggravates the airway pathogenesis in
87 the lungs and necessitates numerous cells and organelle components" (7), according to NIH
88 (National Institutes of Health) recommendations from 1991, 1997 and 2007. Around 339
89 million people worldwide suffer from asthma and associated symptoms in some form. Even
90 industrialized nations are not immune to its effects; thousands of individuals in the United
91 States suffer asthma-like symptoms. In India, it is estimated that one in every three adults
92 suffers from asthma. Coughing fits, wheezing fits, shortness of breath (dyspnea), and chest
93 tightness, are the most likely/common asthma symptoms (8, 9). Many allergens may induce
94 bronchospasm, constriction, respiratory tract infection, hypersensitivity reactions, and
95 contraction of the airway zone, which can make inhalation (or exhalation) difficult. Allergens
96 may also play a role in these responses; however, exercise may also trigger them. Some

97 medications, such as morphine, non-steroidal anti-inflammatory medicines (NSAIDs, such as
98 indomethacin, ibuprofen, fenoprofen, ketorolac, and naproxen), β -antagonists, and angiotensin-
99 converting enzyme (ACE) blockers, for example, perindopril, lisinopril, ramipril, and enalapril
100 (by causing coughing), might exacerbate asthma symptoms (10).

101 Asthma is a systemic condition that starts with the activation of a large number of
102 inflammatory cells, causing lasting alterations in the lower and upper respiratory organs. These
103 modifications in the shape of the bronchial tract are referred to as airway remodeling. Repeated
104 inflammation stimulus *via* matrix peptides and growth factors generated by inflammatory
105 biomolecules and immune cells causes these alterations (11).

106 COVID-19 etiology is more likely to deteriorate in the elderly and those with a medical
107 history of diabetes, severe asthma or heart disease. Although it is thought that people with
108 chronic asthma who are infected with the virus may have poorer results, COVID-19 may
109 increase asthma symptoms in severe asthmatics, much like other viral diseases. Many
110 cytokines, including interleukins (*e.g.*, IL-1, IL-6, IL-12), tumor necrosis factor- α (TNF- α),
111 interferon- γ (IFN- γ) and others, have been shown to be higher in the blood of COVID-19
112 positive individuals, which may have implications for lung inflammation and injury.

113 SARS (severe acute respiratory syndrome) is a kind of acute respiratory syndrome. The
114 heart, blood arteries, lungs, and gut of the host cell are where Coronavirus 2 (SARS-CoV-2)
115 connects to ACE-2 receptors. As a result, COVID-19 is provided to asthmatic patients with
116 significant respiratory symptoms. A sizable fraction of patients has ARDS (acute respiratory
117 distress syndrome), and as a result, the painful symptoms are linked to real hyper cytokinesis
118 specific for IL-6, which may lead to mortality (12).

119 Asthma is most usually caused by respiratory viruses. Among the causes of asthma
120 aggravations, the human rhinovirus is recognized as the most potent and frequent. SARS-CoV-
121 2 is also linked to the progress of asthma, which is a hazard marker for COVID-19 linked injury.
122 COVID-19 causes shortness of breath and a dry cough, and it's also linked to moderate asthma
123 worsening. High temperature is more than often associated with COVID-19, although it may
124 also be detected in some infection-related aggravation of asthma. As a result, COVID-19
125 screening techniques are required for everyone with respiratory issues or asthma (13).

126 As a first-line treatment, severe asthma is managed using inhalers and biological
127 agents/drugs, such as omalizumab, mepolizumab, reslizumab, benralizumab, and dupilumab.
128 Other therapy alternatives, such as bronchial thermoplasty (B.T.), are the two mainstay

129 therapies for severe asthmatics that have not been able to manage their symptoms after
130 receiving first treatment. If any patient has an acute attack, they should contact their physicians
131 if their symptoms worsen. Assume that your severe asthma is under control: there is no need to
132 go to the hospital on a frequent basis in such instances. Patients who meet the emergency
133 requirements must be sent to the hospital right away (14).

134 Researchers from Rutgers University in New Jersey discovered that individuals with no
135 indicators of asthma had nil greater hazard of contracting COVID-19 with respect to asthmatic
136 individuals in research. Researchers identified a drop in T-cell levels in COVID-19, while
137 asthmatic individuals with COVID-19 had higher circulating T-lymphocyte expression, which
138 seems to safeguard them from the severity of SARS-CoV-2 (15).

139

140 *Critical care and management of COVID-19 with asthma*

141 A number of monoclonal antibodies that target type-2 inflammatory pathways and hence,
142 prevent asthma attacks are the most often recommended medications in the treatment of asthma.
143 Omalizumab, anti-IL-4 and anti-IL13 antibodies are found to decrease the worsening of
144 asthmatic symptoms as another kind of therapy. Asthma and COVID-19 have an inflammatory
145 burst trait, making it possible to treat them both with the same treatment. Because of this, anti-
146 inflammatory medicines and biologicals (*e.g.*, monoclonal antibodies) may be used to treat both
147 of these respiratory conditions by targeting a distinct yet common route. Clazakizumab
148 (NCT04343989), tocilizumab (NCT04306705, NCT04346355), and siltuximab
149 (NCT04330638) have all been tested in clinical studies to see how they affect IL-6 in patients
150 (16, 17).

151

152 *Critical care and management of mild cases of COVID-19 with asthma*

153 Patients with asthma who are tested positive for the virus must be quarantined promptly
154 so as to halt the spread of the infection. The affected subject must be isolated in a confined area
155 that has adequate oxygenation. To keep track of the patient's status, a medical checkup must be
156 performed on a regular basis. Local First Referral Units (FRUs), COVID Care Centers, or
157 district/regional community health care centers may treat mild symptoms. The patient's medical
158 history must be documented. Every day, the temperature and oxygen saturation levels must be
159 monitored. A patient's additional risk factors must be taken into consideration. In the event of
160 an emergency, transport the patient to a local COVID hospital. Maintaining a SpO₂ level higher

161 than 94 % is required. Antipyretics such as PCM (paracetamol) should be used in the event of
162 a fever. In order to avoid a dry cough, antitussives may be prescribed. Food and water must be
163 provided to the sufferer (patient) in order for them to recover. If the patient has shortness of
164 breath, he or she should use an emergency inhaler provided by their doctor. Inhaling
165 salbutamol, an alpha-2 agonist, is usually all that is needed to avoid an attack. Prophylactic
166 usage of antimalarial medicine hydroxylchloroquine may be administered to those individuals
167 with hazard issues of more serious illnesses or patients over the age of sixty (averted in case
168 QTc is > 480 ms). Supplemental vitamin C and zinc pills may be used. Hydroxychloroquine
169 may be supplied as a preventative precaution to healthcare workers, such as police and
170 paramilitary, who have a direct touch with patients. On day 1, they may take 400 mg b.d.,
171 trailed by 400 mg one time in 7 days for the next 49 days as the official regimen. Even though
172 children under the age of 15 are not allowed to use it, nursing mothers, and individuals with
173 cardiovascular issues are. Paracetamol is used to treat fever. Favipiravir (favilavir), an anti-
174 influenza medication, was recently licensed for use in India for mild to moderate cases. An
175 RNA polymerase inhibitor having an affinity for RNA viruses is the powerful RNA-dependent
176 RNA polymerase (RNA replication) blocker. In clinical settings, a loading dose of 1.8 g two
177 times on 1st day, which is trailed by a maintenance dose of 0.8 g b.d., and is continued for 7
178 days, is commonly followed (18–20).

179

180 *Critical care and management of moderate cases of COVID-19 with asthma*

181 Patients with mild COVID-19 indicators, such as pneumonia, copious bronchial
182 secretions, and chest blockage, must be kept in isolation at the outset, and if their condition
183 worsens, they should be transported to a neighboring emergency center. The most important
184 clinical parameters, such as respiratory rate and oxygen saturation level (SpO₂), should be
185 checked on a regular basis. A pulse oximeter should always be used to assess oxygen levels. A
186 comprehensive blood count (CBC) accompanied by entire lymphocyte enumeration, renal tests,
187 hepatic markers estimation, 12-lead ECG, X-ray of the thoracic cavity, assessing vital
188 indicators (*e.g.*, breathing, HR), *etc.*, should be performed in order to properly examine the
189 patient. In case SpO₂ goes less than the usual limits, oxygen is administered. PCM 500 mg is a
190 common antipyretic that is used three times daily. If necessary, an antitussive is administered.
191 On the first day, 400 mg of hydroxychloroquine is taken twice a day, trailed by 200 mg two
192 times a day for the next 4 days following the correct ECG check (hydroxychloroquine shall be
193 averted if QTc is > 480 ms). When it comes to treating moderate asthma, beta-2 agonists and

194 steroids such as the combination of salmeterol + fluticasone propionate, or formoterol +
195 budesonide, *etc.*, have been excellent options. Reduced incidence of asthmatic exacerbation has
196 also been shown with the usage of leukotriene receptor antagonists, for instance, montelukast
197 and zafirlukast, as well as 5-lipoxygenase (5-LOX) inhibitors, for instance, zileuton.
198 Methylprednisolone (0.5–1 mg kg⁻¹, intravenously) is given for 3 days in the presence of
199 elevated inflammatory markers. Antibiotics such as azithromycin (500 mg, *p.o.*) once a day and
200 ceftriaxone (1 g, *i.v.*) two times a day, are administered in the co-presence of secondary
201 bacterial infections. Patients exhibiting minor indicators may be offered experimental
202 medications like remdesivir or tocilizumab. Steroids (0.1–0.2 mg kg⁻¹) or methylprednisolone
203 (0.5–1 mg kg⁻¹) are given intravenously for five days (18–20).

204

205 *Critical care and management of severe cases of COVID-19 with asthma*

206 It is necessary to provide intensive treatment to patients with severe COVID-19
207 symptoms and asthma. To treat shock or hypoxemia, such individuals must be supplied oxygen
208 quickly. Oxygen is administered at 5 L min⁻¹ up to SpO₂ is 93–97 %. Children under the age of
209 15 are vented to keep their SpO₂ levels above 94 %. Various kinds of oxygen masks, such as a
210 basic face mask, a nasal cannula, and a mask with a storage-buffer bag, shall every time be
211 available on hand for patients' use. The use of a high-flow nasal cannula (HFNC) for oxygen
212 administration is used if normal oxygen therapy is not being tolerated by patients. HFNC
213 produces 100 % humidified and heated (37 °C) oxygen at a flow rate of 60
214 L min⁻¹. Observation of fluids must be carried out on a frequent basis. Chronic medical
215 monitoring is necessary for patients with ARDS who need daily ventilation for 16–18 hours. In
216 the event of a subsequent bacterial infection, azithromycin 500 mg tablet once in a day and
217 piperacillin/tazobactam (4.5 mg *i.v.*) is administered t.d. for five days. For the first hour, high
218 dosages of inhaled beta-2 agonists are administered, which may be taken for up to four hours.
219 In certain cases, anticoagulants like enoxaparin injection may be necessary. A dosage of 400
220 mg b.d. for one day trailed by 200 mg b.d. for 14 consecutive days is also an option for
221 medications like hydroxychloroquine. Lopinavir 400 mg or ritonavir 100 mg b.d. will be given
222 for the following 14 days. Tocilizumab immunomodulatory treatment may be completed as
223 well if desired. Steroids with dose range 0.2–0.4 mg kg⁻¹ or methylprednisolone in doses 1–2
224 mg kg⁻¹ are administered intravenously over the course of a 10-day period. First-day
225 administration of 200 mg *i.v.* of remdesivir may be trailed by five days of 100 mg *i.v.* for
226 moderate to severe patients. Sputum, CBC, culturing the pleural fluid, and SpO₂ must all be

227 monitored on a regular basis in the event that a patient has pneumonia. Severe acquired
228 pneumonia may be controlled for the following 5 days with doxycycline 200 mg tablet (oral
229 route) on 1stday, trailed by 100 mg (18–20). Table I shows the specific therapy
230 recommendations centered on the seriousness of the COVID-19 indicators.

231

232

SMOKING AND COVID-19

A risk of COVID-19 to smokers

234 More than 1.1 billion people throughout the world smoke every day, and the
235 contemporary figure is expected to soar to 1.3 billion by 2025 as reported by the WHO (21). It
236 is believed that tobacco smoke contains more than 5,000 different compounds, many of which
237 are known carcinogens or poisons (22), and that this combination is the primary source of
238 exposure for both humans and other living species (23). When it comes to the progress and
239 aggravation of several breathing disorders involving microbial invasion as well (24, 25),
240 smoking is a key hazard component. Chronic obstructive pulmonary disease (COPD) and lung
241 melanoma is both clearly linked to tobacco smoking in general (26, 27). It has also been shown
242 that COVID-19 advances more rapidly in COPD patients. Because smoking is a significant
243 element in the progress of COPD, it may have a comparable influence on symptoms (23, 26).
244 Smoking was shown to have no effect on the severity of COVID-19 in a recent meta-analysis
245 (6). The heterogeneity among the studies was modest in this meta-analysis, which comprised
246 just five investigations in total (1).

247 Suffice it to say that smoking has been linked to an expanded hazard of community-
248 acquired pneumonia (CAP) (28, 29) owing to the disruption of respiratory epithelial repair and
249 healing caused by tobacco smoke. One of the most significant risk factors for a respiratory tract
250 infection is smoking, for instance, active, passive and third-hand smoke exposure, all of which
251 include the act of smoking cigarettes (30, 31). Smoking has long been recognized to have a
252 detrimental effect on lung health. Smoking damages the immune strength in the respiratory
253 tract, which renders smokers highly predisposed to undeterred microbial invasions, as
254 previously stated (32). Tobacco smoking has been linked to MERS (Middle East Respiratory
255 Syndrome) infection and death in previous studies (33, 34). There was a greater risk of MERS-
256 related death among smokers compared to non/never-smokers (35). The fact that both MERS-
257 CoV and SARS-CoV-2 are representatives of the Coronaviridae family isn't enough to render

258 people additionally vulnerable to SARS-infection with CoV-2 or a worsening of the prognosis
259 of COVID-19.

260

261 *Critical care and management of COVID-19 patients having smoking history*

262 COVID-19 development is hampered by smoking. In comparison to non-smokers,
263 smokers display a far greater chance of contracting COVID-19 contamination. All communities
264 across the globe have a high prevalence of tobacco smoking (36). An estimated 8–8.5 million
265 people die each year as a result of their dependence on cigarettes. Tobacco smoking exacerbates
266 the issue because COVID-19 spreads by salivary droplets and triggers acute lung pneumonia
267 in humans. Smokers present a bigger probability of COVID-19 illness because of their reduced
268 lung performance and prone cleanliness behaviors (37). Exhaled smoke, coughing, and salivary
269 droplets contaminated with SARS-COV-2 from infected individuals pollute surfaces, posing
270 health risks to those who breathe in this contaminated air (38). The WHO has issued a number
271 of recommendations for preventing the spread of viral illnesses, including washing hands for
272 15–20 seconds with soap and water and using an alcohol-based sanitizer after every encounter.
273 Keep a 1-meter space between you and the next person. Despite this, keep your hands away
274 from your eyes, nose and mouth (39). Tobacco smoking cessation and a reduction might be an
275 effective way to slow the transmission of the virus. Because of this, governments and the
276 general public alike should enact and enforce stringent measures to reduce smoking (40).
277 Smokers should be segregated and only allowed to smoke in designated places that meet strict
278 cleanliness requirements (such as cigarette holders or filters) and have enough ventilation.
279 Conventional cigarettes, electronic cigarettes, and water pipes should not be used for re-use or
280 sharing. After a single usage, cigarette stubs, gadgets, and their attachments should be thrown
281 away to avoid spreading the disease. For those who want to quit smoking, therapeutic nicotine
282 formulations may also be utilized in the process. Nicotine (41) reactivates the nicotine
283 cholinergic system, which had been rendered dysfunctional. Nicotine is widely accepted and
284 has been used for decades due to its low cost and attractive safety compliance (42). On the other
285 hand, since it may be purchased over the counter, there is a risk of abuse (41). As smoking
286 happens to be the single chief reason for severe respiratory disorders such as COPD and
287 emphysema, hence, treatment of these in the ailing patient becomes mandatory even in the
288 presence of COVID-19 symptoms (Table II).

289 *Correlation between angiotensin-converting enzyme (ACE), smoking and COVID-19*

290 In regard to COVID-19, ACE-2 has garnered all the limelight on the planet (43). These
291 include the respiratory system, the heart, and the digestive system (44). ACE-2 is a type II trans-
292 membrane metallo carboxypeptidase subtype that converts angiotensin II into a variety of
293 metabolites, including angiotensin 1–9 and 1–7 (45, 46). Type-II pneumocytes exhibit the
294 presence of ACE-2 (47). It has a critical role in the control of blood pressure and heart activity,
295 however, its involvement in the thoracic cavity is less clear (48, 49). In spite of the fact that
296 certain research has shown no correlation between the ACE-2 genetic polymorphism and
297 COVID-19 invasion, numerous other investigations have found that the receptor plays a major
298 effect (50–53). The structural similarity between the ACE-2 receptor and the coronavirus
299 promotes their interaction (54). Studies demonstrate that smokers reveal an enhanced
300 appearance of the ACE-2 receptor relative to non-smokers (55, 56). SARS-CoV-2 infection
301 may be more likely to occur in smokers because of this (57). New approaches to treating
302 COVID-19, in particular in smokers, may benefit from therapeutic targeting of ACE-2 (Fig. 1).

303

304 *Critical care and management of COVID-19 with asthmatic patient who has smoking history*

305 Tobacco abuses and asthma in COVID-19 patients both have significant negative
306 implications on the patient's overall health, prognosis and treatment options. Compared to non-
307 smokers with asthma, smokers with asthma had a greater risk of developing COVID-19 illness.
308 Smoking in asthma patients with COVID-19 virus alters airway inflammation, ACE-2, and
309 corticosteroid insensitivity, all of which contribute to the negative consequences of smoking.
310 While quitting smoking in a COVID setting may help alleviate symptoms and improve lung
311 function, the poor success rate of this strategy underscores the need for new approaches to the
312 care of these individuals.

313 Many transcriptome data sets of lung samples from healthy never- and ever-smokers and
314 asthmatic patients show that ACE-2 gene expression is up-regulated in smokers relative to non-
315 smokers. ACE-2-producing goblet cells are also found in the lungs of those who have ever
316 smoked compared to those who have never smoked. Many components of the RAS (renin-
317 angiotensin system) are affected by nicotine's action in the body's many organ systems; nicotine
318 up-regulates renin, ACE, and AT1R expression and activity in the ACE/AT-II (angiotensin
319 II)/AT1R 1-7 arm, while down regulating ACE2 and AT2R expression and activity in the
320 compensatory ACE-2/angiotensin 1–7 arm. SARS-CoV-2 may get passage into the lungs more

321 easily *via* the bronchial epithelium of asthmatics who smoke cigarettes, but this does not always
322 mean that they are at greater risk of getting COVID-19 pneumonia.

323

324

CONCLUSIONS

325 The COVID-19 outbreak rocked the globe since it poses a serious danger to human
326 health. Considering the COVID-19 epidemic, severe asthma treatment is now a major issue that
327 will continue until herd immunity is developed soon. Some cytokines have increased in the
328 severe instances of COVID-19, and this might aggravate asthma attacks by causing more
329 inflammation in the bronchial tracts. More research is required, but it seems that biologicals for
330 acute/critical asthma might not raise the possibility of infection or serious COVID-19 and may
331 even be beneficial. Smoking and COVID-19 have been shown to have a direct relationship,
332 with smokers being more susceptible to the infection than non-smokers. Increased ACE-2
333 levels are linked to an enhanced danger of SARS-COV-2 illness in those who smoke.
334 Individuals' susceptibility to the COVID-19 infection may be reduced by treating ACE-2 and
335 quitting smoking. SARS-COV-2 transmission may be slowed down by reducing or quitting
336 smoking. Smoking is a major issue in the current pandemic and cessation of smoking poses a
337 huge challenge to healthcare system across the globe. Smoking cessation must be amongst the
338 primary goals in anti-COVID-19 strategies and also a general health improvement and well-
339 being of people.

340

341 *Abbreviation, acronyms, symbols.* – 2019-nCov – 2019-novel coronavirus, 5-LOX – 5-
342 lipoygenase, ACE – angiotensin-converting enzyme, ARDS – acute respiratory distress syndrome,
343 AT1R – angiotensin II type 1 receptor, AT-II – angiotensin II, b.d. – *bis in die*, B.T. – bronchial
344 thermoplasty, CAP – community-acquired pneumonia, CBC – comprehensive blood count, COPD –
345 chronic obstructive pulmonary disease, COVID-19 – Coronavirus disease 2019, ECG –
346 electrocardiograph, HFNC – high-flow nasal cannula, HR – heart rate, IFN- γ – interferon- γ , IL –
347 interleukin, MERS – Middle East Respiratory Syndrome, NSAIDs – non-steroidal anti-inflammatory
348 drugs, o.d. – *omni die* (every day, once daily), PCM – paracetamol, QTc – QT corrected for heart rate,
349 RNA – ribonucleic acid, SARS – Severe Acute Respiratory Syndrome, SARS-CoV-2 – Severe Acute
350 Respiratory Syndrome Coronavirus 2, SpO₂ – oxygen saturation, t.d. – *ter in die* (three times a day),
351 TNF- α – tumor necrosis factor- α

352 *Acknowledgments.* – We are thankful to Amar Shaheed Baba Ajit Singh Jujhar Singh Memorial
353 College of Pharmacy, Bela (an autonomous college) Ropar (Punjab) to provide the necessary facilities
354 for the completion of this study.

355 *Author's contribution.* – Dongming Lu (conception and design of content and critical revision of
356 the article); Obaid Yaqoob (literature survey and drafting the article); Manish Kumar (English language
357 editing and critical revision of the article); Ajay Singh Kushwah (conception and design of content,
358 write the article; corresponding author); Rahul Kumar Sharma (conceived and designed the content);
359 Devinder Kumar (English language editing and proof reading); Yogendra Mavai (literature survey and
360 critical revision of the article) and Rukaiya Khan (proof reading and language editing).

361 *Conflicts of interest.* – The authors declare no conflicts of interest.

362 *Funding.* – This research has not received any specific grant

363 *ORCIDiDs.* – Dongming Lu – <https://orcid.org/0000-0002-4153-3036>, Obaid Yaqoob –
364 <https://orcid.org/0000-0002-8512-6547>, Manish Kumar – <https://orcid.org/0000-0001-6697-544X>,
365 Ajay Singh Kushwah – <https://orcid.org/0000-0003-0559-7670>, Rahul Kumar Sharma –
366 <https://orcid.org/0000-0002-9678-6882>, Devinder Kumar – <https://orcid.org/0000-0003-2399-408X>,
367 Yogendra Mavai – <https://orcid.org/0000-0002-6241-5047>, Rukaiya Khan – <https://orcid.org/0000-0002-5502-8540>

369

370 REFERENCES

- 371 1. S. Ghosh, S. Das, R. Mondal, S. Abdullah, S. Sultana, S. Singh, A. Sehgal and T. Behl, A review
372 on the effect of COVID-19 in type 2 asthma and its management, *Int. Immunopharmacol.* **91** (2021)
373 Article ID 107309; <https://doi.org/10.1016/j.intimp.2020.107309>
- 374 2. D. Wang, B. Hu, C. Hu, F. Zhu, X. Liu, J. Zhang, B. Wang, H. Xiang, Z. Cheng, Y. Xiong, Y. Zhao,
375 Y. Li, X. Wang and Z. Peng, Clinical characteristics of 138 hospitalized patients with 2019 novel
376 coronavirus-infected pneumonia in Wuhan, China, *JAMA* **323** (2020) 1061–1069;
377 <https://doi.org/10.1001/jama.2020.1585>
- 378 3. O. I. Oyeniran and T. Chia, Novel coronavirus disease 2019 (COVID-19) outbreak in Nigeria: how
379 effective are government interventions?, *Ethics Med. Public Health* **14** (2020) Article ID 100515;
380 <https://doi.org/10.1016/j.jemep.2020.100515>
- 381 4. Q. Li, X. Guan, P. Wu, X. Wang, L. Zhou, Y. Tong, R. Ren, K. S. M. Leung, E. H. Y. Lau, J. Y.
382 Wong, X. Xing, N. Xiang, Y. Wu, C. Li, Q. Chen, D. Li, T. Liu, J. Zhao, M. Liu, W. Tu, C. Chen, L.
383 Jin, R. Yang, Q. Wang, S. Zhou, R. Wang, H. Liu, Y. Luo, Y. Liu, G. Shao, H. Li, Z. Tao, Y. Yang,
384 Z. Deng, B. Liu, Z. Ma, Y. Zhang, G. Shi, T. T. Y. Lam, J. T. Wu, G. F. Gao, B. J. Cowling, B.

- 385 Yang, G. M. Leung and Z. Feng, Early transmission dynamics in Wuhan, China, of novel
386 coronavirus-infected pneumonia, *N. Engl. J. Med.* **382** (2020) 1199–1207;
387 <https://doi.org/10.1056/NEJMoa2001316>
- 388 5. J. Hartmann-Boyce, J. Gunnell, J. Drake, A. Otunla, J. Suklan, E. Schofield, J. Kinton, M. Inada-
389 Kim, F. D. R. Hobbs and P. Dennison, Asthma and COVID-19: review of evidence on risks and
390 management considerations, *BMJ Evid-Based Med.* **26**(4) (2021) Article ID 195(8 pages);
391 <https://doi.org/10.1136/bmjebm-2020-111506>
- 392 6. I. B. Ahluwalia, M. Myers and J. E. Cohen, COVID-19 pandemic: an opportunity for tobacco use
393 cessation, *Lancet Pub. Health* **5** (2020) e577; [https://doi.org/10.1016/S2468-2667\(20\)30236-X](https://doi.org/10.1016/S2468-2667(20)30236-X)
- 394 7. J. W. Mims, Asthma: definitions and pathophysiology, *Int. Forum Allergy Rhinol.* **5** (2015) S2–S6;
395 <https://doi.org/10.1002/alr.21609>
- 396 8. J. A. Krishnan, R. F. Lemanske, G. J. Canino, K. S. Elward, M. Kattan, E. C. Matsui, H. Mitchell,
397 E. R. Sutherland and M. Minnicozzi, Asthma outcomes: Symptoms, *J. Allergy Clin. Immunol.* **129**
398 (2012) S124–S135; <https://doi.org/10.1016/j.jaci.2011.12.981>
- 399 9. J. Bousquet, E. Mantzouranis, A. A. Cruz, N. Ait-Khaled, C. E. Baena-Cagnani, E. R. Bleeker, C.
400 E. Brightling, P. Burney, A. Bush, W. W. Busse, T. B. Casale, M. Chan-Yeung, R. Chen, B.
401 Chowdhury, K. F. Chung, R. Dahl, J. M. Drazen, L. M. Fabbri, S. T. Holgate, F. Kauffmann, T.
402 Haahtela, N. Khaltaev, J. P. Kiley, M. R. Masjedi, Y. Mohammad, P. O'Byrne, M. R. Partridge, K.
403 F. Rabe, A. Togias, C. van Weel, S. Wenzel, N. Zhong and T. Zuberbier, Uniform definition of
404 asthma severity, control, and exacerbations: Document presented for the World Health Organization
405 consultation on severe asthma, *J. Allergy Clin. Immunol.* **126** (2010) 926–938;
406 <https://doi.org/10.1016/j.jaci.2010.07.019>
- 407 10. F. T. Ishmael, The inflammatory response in the pathogenesis of asthma, *J. Am. Osteopath. Assoc.*
408 **111**(11, Suppl. 7) (2011) S11–S17; <https://doi.org/10.7556/jaoa.2011.20014>
- 409 11. C. J. Oliphant, J. L. Barlow and A. N. J. McKenzie, Insights into the initiation of type 2 immune
410 responses, *Immunology* **134** (2011) 378–385; <https://doi.org/10.1111/j.1365-2567.2011.03499.x>
- 411 12. M. Morais-Almeida and J. Bousquet, COVID-19 and asthma: To have or not to have T2
412 inflammation makes a difference?, *Pulmonology* **26**(5) (2020) 261–263;
413 <https://doi.org/10.1016/j.pulmoe.2020.05.003>
- 414 13. E. M. Abrams, G. W. 't Jong and C. L. Yang, Asthma and COVID-19, *CMAJ* **192**(20) (2020) E551;
415 <https://doi.org/10.1503/cmaj.200617>
- 416 14. T. Tabassum, A. Rahman, Y. Araf, Md. A. Ullah and M. J. Hosen, Management of asthma patients
417 during the COVID-19 pandemic: pathophysiological considerations to address the challenges, *Beni-*

- 418 *Suef Univ. J. Basic Appl. Sci.* **11** (2022) Article ID 20(14 pages); [https://doi.org/10.1186/s43088-](https://doi.org/10.1186/s43088-022-00204-4)
419 022-00204-4
- 420 15. K. E. J. Philip, S. Buttery, P. Williams, B. Vijayakumar, J. Tonkin, A. Cumella, L. Renwick, L.
421 Ogden, J. K. Quint, S. L. Johnston, M. I. Polkey and N. S. Hopkinson, Impact of COVID-19 on
422 people with asthma: a mixed methods analysis from a UK wide survey, *BMJ Open Resp. Res.* **9**
423 (2022) e001056; <https://doi.org/10.1136/bmjresp-2021-001056>
- 424 16. J. Darveaux and W. W. Busse, Biologics in asthma – The next step toward personalized treatment,
425 *J. Allergy Clin. Immunol.: In Practice* **3** (2015) 152–160; <https://doi.org/10.1016/j.jaip.2014.09.014>
- 426 17. S. Atal and Z. Fatima, IL-6 Inhibitors in the treatment of serious COVID-19: A promising therapy?,
427 *Pharm. Med.* **34** (2020) 223–231; <https://doi.org/10.1007/s40290-020-00342-z>
- 428 18. U. Agrawal, R. Raju and Z. F. Udawadia, Favipiravir: A new and emerging antiviral option in
429 COVID-19, *Med. J. Armed Forces India* **76**(4) (2020) 370–376;
430 <https://doi.org/10.1016/j.mjafi.2020.08.004>
- 431 19. S. Sharma, S. Basu, N. P. Shetti and T. M. Aminabhavi, Current treatment protocol for COVID-19
432 in India, *Sensors Int.* **1** (2020) Article ID 100013(3 pages);
433 <https://doi.org/10.1016/j.sintl.2020.100013>
- 434 20. A. Parasher, COVID-19: Current understanding of its pathophysiology, clinical presentation and
435 treatment, *Postgrad. Med. J.* **97** (1147) (2021) 312–320; [https://doi.org/10.1136/postgradmedj-](https://doi.org/10.1136/postgradmedj-2020-138577)
436 2020-138577
- 437 21. V. Bilano, S. Gilmour, T. Moffiet, E. T. d’Espaignet, G. A. Stevens, A. Commar, F. Tuyt, I. Hudson
438 and K. Shibuya, Global trends and projections for tobacco use, 1990–2025: an analysis of smoking
439 indicators from the WHO comprehensive information systems for tobacco control, *Lancet* **385**
440 (2015) 966–976; [https://doi.org/10.1016/s0140-6736\(15\)60264-1](https://doi.org/10.1016/s0140-6736(15)60264-1)
- 441 22. R. Talhout, T. Schulz, E. Florek, J. Van Benthem, P. Wester and A. Opperhuizen, Hazardous
442 compounds in tobacco smoke, *Int. J. Environ. Res. Public Health* **8** (2011) 613–628;
443 <https://doi.org/10.3390/ijerph8020613>
- 444 23. M. Ezzati and A. D. Lopez, Estimates of global mortality attributable to smoking in 2000, *Lancet*
445 **362** (2003) 847–852; [https://doi.org/10.1016/s0140-6736\(03\)14338-3](https://doi.org/10.1016/s0140-6736(03)14338-3)
- 446 24. S. J. Brake, K. Barnsley, W. Lu, K. D. McAlinden, M. S. Eapen and S. S. Sohal, Smoking
447 upregulates angiotensin-converting enzyme-2 receptor: A potential adhesion site for novel
448 coronavirus SARS-CoV-2 (Covid-19), *J. Clin. Med.* **9**(3) (2020) Article ID 841(7 pages);
449 <https://doi.org/10.3390/jcm9030841>
- 450 25. M. D. Shastri, S. D. Shukla, W. C. Chong, Rajendra KC, K. Dua, R. P. Patel, G. M. Peterson and

- 451 R. F. O'Toole, Smoking and COVID-19: What we know so far, *Resp. Med.* **176** (2021) Article ID
452 106237 (7 pages); <https://doi.org/10.1016/j.rmed.2020.106237>
- 453 26. N. M. Siafakas, P. Vermeire, N. B. Pride, P. Paoletti, J. Gibson, P. Howard, J. C. Yernault, M.
454 Decramer, T. Higenbottam, D. S. Postma and J. Rees, Optimal assessment and management of
455 chronic obstructive pulmonary disease (COPD), *Eur. Resp. J.* **8** (1995) 1398–1420;
456 <https://doi.org/10.1183/09031936.95.08081398>
- 457 27. D. Milner, The physiological effects of smoking on the respiratory system, *Nurs. Times* **100** (2004)
458 56–59.
- 459 28. J. Almirall, I. Bolibar, M. Serra-Prat, J. Roig, I. Hospital, E. Carandell, M. Agusti, P. Ayuso, A.
460 Estela and A. Torres, New evidence of risk factors for community-acquired pneumonia: a
461 population-based study, *Eur. Resp. J.* **31** (2008) 1274–1284;
462 <https://doi.org/10.1183/09031936.00095807>
- 463 29. H. Müllerova, C. Chigbo, G. W. Hagan, M. A. Woodhead, M. Miravitlles, K. J. Davis and J. A.
464 Wedzicha, The natural history of community-acquired pneumonia in COPD patients: A population
465 database analysis, *Resp. Med.* **106** (2012) 1124–1133; <https://doi.org/10.1016/j.rmed.2012.04.008>
- 466 30. S. Cohen, D. A. Tyrrell, M. A. Russell, M. J. Jarvis and A. P. Smith, Smoking, alcohol consumption,
467 and susceptibility to the common cold, *Am. J. Public Health* **83** (1993) 1277–1283;
468 <https://doi.org/10.2105/ajph.83.9.1277>
- 469 31. J. D. Kark, M. Lebiush and L. Rannon, Cigarette smoking as a risk factor for epidemic A(H1N1)
470 influenza in young men, *N. Engl. J. Med.* **307** (1982) 1042–1046;
471 <https://doi.org/10.1056/nejm198210213071702>
- 472 32. Z. Zhou, P. Chen and H. Peng, Are healthy smokers really healthy?, *Tob. Ind. Dis.* **14** (2016) Article
473 ID 35 (12 pages); <https://doi.org/10.1186/s12971-016-0101-z>
- 474 33. L. Arcavi and N. L. Benowitz, Cigarette smoking and infection, *Arch. Intern. Med.* **164** (2004)
475 2206–2216; <https://doi.org/10.1001/archinte.164.20.2206>
- 476 34. J.-E. Park, S. Jung, A. Kim and J.-E. Park, MERS transmission and risk factors: a systematic review,
477 *BMC Public Health* **18** (2018) Article ID 574 (15 pages); [https://doi.org/10.1186/s12889-018-5484-](https://doi.org/10.1186/s12889-018-5484-8)
478 8
- 479 35. H.-S. Nam, J. W. Park, M. Ki, M.-Y. Yeon, J. Kim and S. W. Kim, High fatality rates and associated
480 factors in two hospital outbreaks of MERS in Daejeon, the Republic of Korea, *Int. J. Infect. Dis.* **58**
481 (2017) 37–42; <https://doi.org/10.1016/j.ijid.2017.02.008>
- 482 36. C. Sherman, Health effects of cigarette smoking, *Clin. Chest Med.* **12** (1991) 643–658;
483 [https://doi.org/10.1016/S0272-5231\(21\)00814-5](https://doi.org/10.1016/S0272-5231(21)00814-5)

- 484 37. K. Palipudi, S. A. Rizwan, D. N. Sinha, L. J. Andes, R. Amarchand, A. Krishnan and S. Asma
485 Prevalence and sociodemographic determinants of tobacco use in four countries of the World Health
486 Organization: South-East Asia region: findings from the global adult tobacco survey, *Indian J.*
487 *Cancer* **51** (Suppl. 1) (2014) S24–S32; [https://doi.org/ 10.4103/0019-509X.147446](https://doi.org/10.4103/0019-509X.147446)
- 488 38. R. M. Benjamin, Exposure to tobacco smoke causes immediate damage: A report of the surgeon
489 general, *Public Health Rep.* **126** (2011) 158–159; <https://doi.org/10.1177/003335491112600202>
- 490 39. W.-J. Guan, Z.-Y. Ni, Y. Hu, W.-H. Liang, C.-Q. Ou, J.-X. He, L. Liu, H. Shan, C.-L. Lei, D. S.C.
491 Hui, B. Du, L.-J. Li, G. Zeng, K.-Y. Yuen, R.-C. Chen, C.-L. Tang, T. Wang, P.-Y. Chen, J. Xiang,
492 S.-Y. Li, J.-L. Wang, Z.-J. Liang, Y.-X. Peng, L. Wei, Y. Liu, Y.-H. Hu, P. Peng, J.-M. Wang, J.-
493 Y. Liu, Z. Chen, G. Li, Z.-J. Zheng, S.-Q. Qiu, J. Luo, C.-J. Ye, S.-Y. Zhu and N.-S. Zhong (for the
494 China Medical Treatment Expert Group for Covid-19), Clinical characteristics of coronavirus
495 disease 2019 in China, *N. Engl. J. Med.* **382** (2020) 1708–1720;
496 <https://doi.org/10.1056/nejmoa2002032>
- 497 40. K. Vapalahti, A.-M. Virtala, A. Vaehri and O. Vapalahti, Case-control study on Puumala virus
498 infection: smoking is a risk factor, *Epidemiol. Infect.* **138** (2009) 576–584;
499 <https://doi.org/10.1017/s095026880999077x>
- 500 41. A. K. Boutou, G. Pitsiou, T. Kontakiotis and I. Kioumis, Nicotine treatment and smoking cessation
501 in the era of COVID-19 pandemic: an interesting alliance, *ERJ Open Res.* **6**(3) (2020) ID 00306-
502 2020(3 pages); <https://doi.org/10.1183/23120541.00306-2020>
- 503 42. H. A. Tindle, P. A. Newhouse and M. S. Freiberg, Beyond smoking cessation: Investigating
504 medicinal nicotine to prevent and treat COVID-19, *Nicotine Tob. Res.* **22**(9) (2020) 1669–1670;
505 <https://doi.org/10.1093/ntr/ntaa077>
- 506 43. S. J. Brake, K. Barnsley, W. Lu, K. D. McAlinden, M. S. Eapen and S. S. Sohal, Smoking
507 upregulates angiotensin-converting enzyme-2 receptor: A potential adhesion site for novel
508 coronavirus SARS-CoV-2 (Covid-19), *J. Clin. Med.* **9**(3) (2020) Article ID 841 (7 pages);
509 <https://doi.org/10.3390/jcm9030841>
- 510 44. K. P. F. Chan, T. F. Ma, W. C. Kwok, J. K. C. Leung, K. Y. Chiang, J. C. M. Ho, D. C. L. Lam, T.
511 C. C. Tam, M. S. M. Ip and P. L. Ho, Significant reduction in hospital admissions for acute
512 exacerbation of chronic obstructive pulmonary disease in Hong Kong during coronavirus disease
513 2019 pandemic, *Resp. Med.* **171** (2020) Article ID 106085 (6 pages);
514 <https://doi.org/10.1016/j.rmed.2020.106085>
- 515 45. M. A. Crackower, R. Sarao, G. Y. Oudit, C. Yagil, I. Kozieradzki, S. E. Scanga, A. J. Oliveira-dos-
516 Santos, J. da Costa, L. Zhang, Y. Pei, J. Scholey, C. M. Ferrario, A. S. Manoukian, M. C. Chappell,
517 P. H. Backx, Y. Yagil and J. M. Penninger, Angiotensin-converting enzyme 2 is an essential

- 518 regulator of heart function, *Nature* **417** (2002) 822–828; <https://doi.org/10.1038/nature00786>
- 519 46. D. Harmer, M. Gilbert, R. Borman and K. L. Clark, Quantitative mRNA expression profiling of
520 ACE 2, a novel homologue of angiotensin converting enzyme, *FEBS Lett.* **532** (2002) 107–110;
521 [https://doi.org/10.1016/s0014-5793\(02\)03640-2](https://doi.org/10.1016/s0014-5793(02)03640-2)
- 522 47. I. Hamming, W. Timens, M. Bulthuis, A. Lely, G. Navis and H. van Goor, Tissue distribution of
523 ACE2 protein, the functional receptor for SARS coronavirus. A first step in understanding SARS
524 pathogenesis, *J. Pathol.* **203** (2004) 631–637; <https://doi.org/10.1002/path.1570>
- 525 48. S. D. Crowley, S. B. Gurley, M. I. Oliverio, A. K. Pazmino, R. Griffiths, P. J. Flannery, R. F.
526 Spurney, H.-S. Kim, O. Smithies, T. H. Le and T. M. Coffman, Distinct roles for the kidney and
527 systemic tissues in blood pressure regulation by the renin-angiotensin system, *J. Clin. Invest.* **115**
528 (2005) 1092–1099; <https://doi.org/10.1172/JCI23378>
- 529 49. C. Tikellis and M. C. Thomas, Angiotensin-converting enzyme 2 (ACE2) is a key modulator of the
530 renin angiotensin system in health and disease, *Int. J. Pept.* **2012** (2012) Article ID 256294 (8
531 pages); <https://doi.org/10.1155/2012/256294>
- 532 50. B. Coutard, C. Valle, X. de Lamballerie, B. Canard, N. G. Seidah and E. Decroly, The spike
533 glycoprotein of the new coronavirus 2019-nCoV contains a furin-like cleavage site absent in CoV
534 of the same clade, *Antiviral Res.* **176** (2020) Article ID 104742 (5 pages);
535 <https://doi.org/10.1016/j.antiviral.2020.104742>
- 536 51. K. E. Follis, J. York and J. H. Nunberg, Furin cleavage of the SARS coronavirus spike glycoprotein
537 enhances cell–cell fusion but does not affect virion entry, *Virology* **350** (2006) 358–369;
538 <https://doi.org/10.1016/j.virol.2006.02.003>
- 539 52. F. Li, Structure of SARS coronavirus spike receptor-binding domain complexed with receptor,
540 *Science* **309** (2005) 1864–1868; <https://doi.org/10.1126/science.1116480>
- 541 53. M. Saadat, No significant correlation between ACE Ins/Del genetic polymorphism and COVID-19
542 infection, *Clin. Chem. Lab. Med.* **58** (2020) 1127–1128; <https://doi.org/10.1515/cclm-2020-0577>
- 543 54. D. Wrapp, N. Wang, K. S. Corbett, J. A. Goldsmith, C.-L. Hsieh, O. Abiona, B. S. Graham and J.
544 S. McLellan, Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation, *Science*
545 **367** (2020) 1260–1263; <https://doi.org/10.1126/science.abb2507>
- 546 55. J. M. Leung, C. X. Yang, A. Tam, T. Shaipanich, T.-L. Hackett, G. K. Singhera, D. R. Dorscheid
547 and D. D. Sin, ACE-2 expression in the small airway epithelia of smokers and COPD patients:
548 Implications for COVID-19, *Eur. Respir. J.* **55**(5) (2020) Article ID 2000688 (5 pages);
549 <https://doi.org/10.1183/13993003.00688-2020>
- 550 56. J. C. Smith, E. L. Sausville, V. Girish, M. L. Yuan, K. M. John and J. M. Sheltzer, Cigarette smoke

551 exposure and inflammatory signaling increase the expression of the SARS-CoV-2 receptor ACE2
552 in the respiratory tract, *Devel. Cell* **53** (2020) 514–529; bioRxiv preprint posted April 26, 2020;
553 <https://doi.org/10.1101/2020.03.28.013672>;

554 57. J. Qi, Y. Zhou, J. Hua, L. Zhang, J. Bian, B. Liu, Z. Zhao and S. Jin, The scRNA-seq expression
555 profiling of the receptor ACE2 and the cellular protease TMPRSS2 reveals human organs
556 susceptible to SARS-CoV-2 infection, *Int. J. Environ. Res. Public Health* **18**(1) (2021) Article ID
557 284 (17 pages); <https://doi.org/10.3390/ijerph18010284>

558

Uncorrected proofs

Table I. Basic care and treatment options of Covid-19 positive asthmatic patients


Mild cases	Reference(s)
<ul style="list-style-type: none"> • Attack aversion by β_2 agonists (<i>e.g.</i>, salbutamol <i>via</i> nasal route) 	1
<ul style="list-style-type: none"> • Multivitamin supplements, for example, vitamin C, D, E and zinc 	1, 12
<ul style="list-style-type: none"> • Anti-tussive, <i>e.g.</i>, dextromethorphan, benzonatate 	1, 14
<ul style="list-style-type: none"> • Persons working in healthcare: As prophylactic hydroxychloroquine (400 mg two times on 1st day upto 49 days with 400 mg once every 7 days (contraindicated < 15 years age, breastfeeding mothers, and cardiovascular patients)) 	1
<ul style="list-style-type: none"> • Antipyretics (<i>e.g.</i>, paracetamol) in high body temperatures 	13
<ul style="list-style-type: none"> • Anti-influenza therapy (<i>e.g.</i>, favipiravir 1.8 g b.d. on 1st day, 0.8 g b.d. for 14 days) 	1
Moderate cases	
<ul style="list-style-type: none"> • If SpO₂ < 92% then O₂ provided 	13
<ul style="list-style-type: none"> • Hydroxychloroquine (400 mg b.d. on 1st day, 200 mg b.d. 2nd day onwards up to 6th day) 	14
<ul style="list-style-type: none"> • Multivitamins and supplements, <i>e.g.</i>, vitamin C and zinc tablets 	18
<ul style="list-style-type: none"> • Anti-tussive, <i>e.g.</i>, dextromethorphan, benzonatate 	1
<ul style="list-style-type: none"> • Avert asthma (β_2 agonists +corticosteroids,<i>e.g.</i>, salmeterol + fluticasone propionate, or formoterol + budesonide, <i>etc.</i>) 	14
<ul style="list-style-type: none"> • Antipyretics (paracetamol) for high body temperature 	1
<ul style="list-style-type: none"> • Abrogate asthmatic aggravation (leukotriene receptor antagonist,<i>e.g.</i>, montelukast, zafirlukast and 5-lipoxygenase inhibitor, <i>e.g.</i>, zileuton) 	16
<ul style="list-style-type: none"> • Methylprednisolone (0.5–1 mg kg⁻¹ day 1–3, <i>i.v.</i>) to abolish inflammatory cascade 	14
<ul style="list-style-type: none"> • Azithromycin (500 mg o.d. for 5 days), ceftriaxone (1000 mg, <i>i.v.</i> b.d. for 5 days for bacterial infection 	18
<ul style="list-style-type: none"> • Remdesivir or tocilizumab in moderate symptoms as investigational drugs 	19
Severe cases	
<ul style="list-style-type: none"> • O₂ 5 L min⁻¹ till SpO₂ is > 92–96 % 	1, 20
<ul style="list-style-type: none"> • Children < 15 years of age are aerated to retain SpO₂ > 94 % 	1
<ul style="list-style-type: none"> • Vitamin C and zinc tablets 	1
<ul style="list-style-type: none"> • Anti-tussive, <i>e.g.</i>, dextromethorphan, benzonatate 	1, 12
<ul style="list-style-type: none"> • High β_2 agonists doses after 15–20 min to 240 min 	1, 13
<ul style="list-style-type: none"> • Hydroxychloroquine (400 mg b.d. on 1st day, 400 mg b.d. 2nd day onwards up to 14th day) 	19
<ul style="list-style-type: none"> • Lopinavir or ritonavir (100 mg b.d.) against viral infection 	18
<ul style="list-style-type: none"> • Antipyretics (paracetamol) in high body temperature 	13
<ul style="list-style-type: none"> • Remdesivir (200 mg <i>i.v.</i>, 1st day and 100 mg 2nd – 5th day) 	19
<ul style="list-style-type: none"> • Doxycycline oral 200 mg tablet on 1st day and 100 mg from 2nd – 6th day (for severe acquired pneumonia) 	1, 13, 20
<ul style="list-style-type: none"> • Azithromycin (500 mg o.d. for 5 days), piperacillin/tazobactam (4.5 mg <i>i.v.</i> t.d.for 5 days) 	19
<ul style="list-style-type: none"> • Dexamethasone (0.2–0.4 mg kg⁻¹) or methylprednisolone (1–2 mg kg⁻¹) for 10 days 	18

Uncorrected proofs

Table II. Therapeutic options of Covid-19 positive smoker patients

Therapy	Reference(s)
Salbutamol (β_2 agonists): 1–2 puffs if breathless occurs up to a maximum of 4–5 times a day ^a	14, 19, 25
Formoterol (β_2 agonists): 12 μg (inhalation) every 12-hour maximum dose 24 μg	19, 26
Indacaterol (β_2 agonists): 75 μg orally once a day	26
Terbutaline (β_2 agonists): 250 μg , 3 to 4 times per day by nebulizer	13
Ipratropium (antimuscarinic): 500 mg <i>via</i> nebulizer 3 or 4 times per day	13, 14, 26
Tiotropium (antimuscarinic): 2.5 μg , maximum dose 2 puffs per day	14, 25
Glycopyrronium (antimuscarinic): 25 μg , <i>i.e.</i> , one inhalation b.d.	14, 26
Aclidinium (antimuscarinic): 400 μg , <i>i.e.</i> , one inhalation b.d.	1
Beclomethasone dipropionate: 800 μg daily	14, 25
Budesonide: 200 μg , 2 inhalations orally twice a day, max dose is 4 inhalations	1
Theophylline: Not more than 400 mg per day	1
Carbocisteine syrup (mucolytic): 15 mL t.d.	14
Carbocisteine capsule (mucolytic): 375 mg t.d.	13, 14
Doxycycline: 100 mg daily	19, 20
Azithromycin: 250–500 mg daily for 1 week	18, 20
Roflumilast: 250 μg per day for 28 days, max dose is 500 μg per day for 28 days	19, 20
O ₂ therapy: 16 hours a day or as required	1, 19
Nicotine replacement therapy	41
Varenicline (reduce craving for nicotine)	41
Antipyretics (<i>e.g.</i> , paracetamol) in high body temperatures	18, 19
Anti-influenza therapy (<i>e.g.</i> , favipiravir 1.8 g b.d. on 1 st day, 0.8 g b.d. for 14 days)	1, 18-20
Remdesivir or tocilizumab	19, 20

563 ^a1 puff = 100 μg



564

565 Fig. 1. Relation between SARS-CoV 2 and ACE-2 receptor in smoking patients.

566

567